

Fermionic functionals without Grassmann numbers

Hrvoje Nikolić

*Theoretical Physics Division, Rudjer Bošković Institute, P.O.B. 180, HR-10002 Zagreb, Croatia.**

(Dated: February 1, 2008)

Since any fermionic operator ψ can be written as $\psi = q + ip$, where q and p are hermitian operators, we use the eigenvalues of q and p to construct a functional formalism for calculating matrix elements that involve fermionic fields. The formalism is similar to that for bosonic fields and does not involve Grassmann numbers. This makes possible to perform numerical fermionic lattice computations that are much faster than not only other algorithms for fermions, but also algorithms for bosons.

PACS numbers: 03.65.Db, 11.15.Tk

Since $\hat{\psi}^2 = 0$ for a fermionic field operator $\hat{\psi}$, this operator does not possess nonzero c-number eigenvalues. Therefore, the functional formalism for fermions involves anticommuting Grassmann numbers [1]. This makes the notion of a classical fermion field conceptually difficult. This is also problematic from a practical point of view, because the functional integrals over Grassmann numbers cannot be performed numerically on a computer, which makes lattice calculations with dynamical fermions difficult. For typical Hamiltonians, the integration over fermionic fields can be reduced to the computation of the determinants of huge matrices, which costs a lot of computer time. Therefore, most of the existing lattice calculations are based on a quenched approximation, in which fermions are not treated dynamically (see, e.g., [2] and references therein). Various methods to improve and speed up the treatment of fermions on a lattice have been proposed, such as those in Refs. [3, 4, 5, 6, 7]. (Ref. [7] contains a longer list of references for various methods.) However, the efficiency of all these existing methods cannot be compared with the efficiency of methods for functional integration over bosonic fields [8].

In this Letter, we construct an entirely new method of functional calculus for fermionic fields. It is based on the fact that any linear operator $\hat{\psi}$ can be written in a unique way as a sum of a hermitian and an antihermitian operator. In other words, it can be written as $\hat{\psi} = \hat{q} + i\hat{p}$, where $\hat{q}$ and $\hat{p}$ are hermitian operators. Since hermitian operators can always be diagonalized, we can use the eigenvalues and eigenstates of $\hat{q}$ and $\hat{p}$ to construct a functional formalism similar to that for bosons. We shall see that $\hat{q}$ and $\hat{p}$ play a role similar to that of the canonical coordinate and the canonical momentum for bosons. This also suggests that $\hat{q}$ and $\hat{p}$ can be naturally interpreted as observables for fermionic fields, which may be relevant for a deeper understanding of the physical meaning of quantum and classical fermionic fields. However, we emphasize the practical utility of our method because the functional integration over fermionic degrees of freedom is not only very similar to that for bosons, but the numerical computation is even much faster than that for bosons. This is because, for each bosonic degree of freedom, the eigenvalues q and p can take any real

value, so lattice calculations require Monte-Carlo simulations with a large number of different real values. On the other hand, it appears that, for each fermionic degree of freedom, q and p can take only two different values.

First, consider one fermionic degree of freedom described by the operators $\hat{\psi}$ and $\hat{\psi}^\dagger$ that satisfy the anticommutation relations

$$\{\hat{\psi}, \hat{\psi}\} = \{\hat{\psi}^\dagger, \hat{\psi}^\dagger\} = 0, \quad \{\hat{\psi}, \hat{\psi}^\dagger\} = 1. \quad (1)$$

This algebra can be represented on a 2-dimensional unitary space spanned by the orthonormal basis vectors $|0\rangle$ and $|1\rangle$ that satisfy $\hat{\psi}|0\rangle = 0$, $\hat{\psi}^\dagger|1\rangle = 0$, $\hat{\psi}^\dagger|0\rangle = |1\rangle$, and $\hat{\psi}|1\rangle = |0\rangle$. The operators $\hat{\psi}$ and $\hat{\psi}^\dagger$ can be written as

$$\hat{\psi} = \hat{q} + i\hat{p}, \quad \hat{\psi}^\dagger = \hat{q} - i\hat{p}, \quad (2)$$

where

$$\hat{q} = (\hat{\psi}^\dagger + \hat{\psi})/2, \quad \hat{p} = i(\hat{\psi}^\dagger - \hat{\psi})/2. \quad (3)$$

are hermitian operators. They satisfy the anticommutation relations

$$\{\hat{q}, \hat{q}\} = \{\hat{p}, \hat{p}\} = 1/2, \quad \{\hat{q}, \hat{p}\} = 0. \quad (4)$$

In particular, we see that $\hat{q}^2 = \hat{p}^2 = 1/4$, which implies that the only eigenvalues of $\hat{q}$ and $\hat{p}$ are $q_\pm = \pm 1/2$ and $p_\pm = \pm 1/2$, respectively. The corresponding eigenvectors are

$$|q_\pm\rangle = (|0\rangle \pm |1\rangle)/\sqrt{2}, \quad |p_\pm\rangle = (|0\rangle \pm i|1\rangle)/\sqrt{2}, \quad (5)$$

with the properties $\hat{q}|q\rangle = q|q\rangle$, $\hat{p}|p\rangle = p|p\rangle$, for q (p) equal to q_+ or q_- (p_+ or p_-). These eigenvectors satisfy the orthonormality relations

$$\langle q|q'\rangle = \delta_{qq'}, \quad \langle p|p'\rangle = \delta_{pp'}, \quad (6)$$

and the completeness relations

$$\sum_q |q\rangle\langle q| = 1, \quad \sum_p |p\rangle\langle p| = 1. \quad (7)$$

They also satisfy

$$\langle p|q\rangle = \frac{1}{\sqrt{2}}(1 - i4pq) = \frac{1}{\sqrt{2}}e^{-i\pi pq}, \quad (8)$$

which is similar to the analogous equation for bosons. (In order to obtain even closer similarity to bosons, one can remove the number π in the last expression in (8) by an appropriate rescaling of the operators $\hat{q}$, $\hat{p}$, and their eigenvalues.)

Let us now apply the above equations to a calculation of the partition function $Z = \text{Tr} \exp(-\beta \hat{H})$, where $\hat{H} = \omega \hat{\psi}^\dagger \hat{\psi}$ is the Hamiltonian and $T = 1/\beta$ is the temperature. Since $\hat{\psi}^\dagger \hat{\psi} |n\rangle = n |n\rangle$ for n equal to 0 or 1, the simplest way to calculate Z is as follows [9]: $Z = \sum_n \langle n | \exp(-\beta \hat{H}) | n \rangle = \exp(-\beta \omega 0) + \exp(-\beta \omega 1) = 1 + \exp(-\beta \omega)$. However, this simple method does not work for a more general case with many degrees of free-

dom and complicated interactions. Therefore, we present a different method, completely analogous to the functional method for bosons [9], that can be generalized to more general cases. We first write the partition function as

$$Z = \sum_q \langle q | e^{-\beta \hat{H}} | q \rangle = \sum_q \langle q | \left(\prod_{j=1}^N e^{-\Delta\tau \hat{H}} \right) | q \rangle, \quad (9)$$

where $\Delta\tau = \beta/N$. The next step is to insert the factor 1 in the forms of (7) $2N$ times and to put N to infinity, so as to obtain

$$Z = \lim_{N \rightarrow \infty} \sum_q \left(\prod_{j=1}^N \sum_{q(j)} \sum_{p(j)} \right) \langle q | p(N) \rangle \langle p(N) | e^{-\Delta\tau \hat{H}} | q(N) \rangle \langle q(N) | p(N-1) \rangle \\ \times \langle p(N-1) | e^{-\Delta\tau \hat{H}} | q(N-1) \rangle \cdots \langle q(2) | p(1) \rangle \langle p(1) | e^{-\Delta\tau \hat{H}} | q(1) \rangle \langle q(1) | q \rangle. \quad (10)$$

Since the Hamiltonian is $\hat{H} = \omega \hat{\psi}^\dagger \hat{\psi} = \omega(1/2 - 2i\hat{p}\hat{q}) \equiv H(\hat{p}, \hat{q})$ and since $\Delta\tau \rightarrow 0$, we can write

$$\langle p(j) | e^{-\Delta\tau H(\hat{p}, \hat{q})} | q(j) \rangle = \langle p(j) | 1 - \Delta\tau H(\hat{p}, \hat{q}) | q(j) \rangle = \langle p(j) | q(j) \rangle e^{-\Delta\tau H(p(j), q(j))}. \quad (11)$$

Therefore, using (11), (8), and (6), (10) can be written as

$$Z = \lim_{N \rightarrow \infty} \sum_q \left(\prod_{j=1}^N \frac{1}{2} \sum_{q(j)} \sum_{p(j)} \right) \delta_{q(1), q} e^{-S}, \quad (12)$$

where

$$S = \Delta\tau \sum_{j=1}^N \left[-i\pi \frac{q(j+1) - q(j)}{\Delta\tau} p(j) + H(p(j), q(j)) \right], \quad (13)$$

and $q(N+1) \equiv q$. The summation over q is easily performed, so the final expression is

$$Z = \lim_{N \rightarrow \infty} \left(\prod_{j=1}^N \frac{1}{2} \sum_{q(j)} \sum_{p(j)} \right) e^{-S}, \quad (14)$$

where the periodic boundary condition $q(N+1) = q(1)$ is understood. Similarity to the bosonic case [9] is obvious. As we have already discussed, the factor π in (13) can be eliminated by an appropriate rescaling. The only important difference is the fact that for bosons one has to make the replacement $(1/2) \sum_{q(j)} \sum_{p(j)} \rightarrow (1/2\pi) \int dq(j) \int dp(j)$. Since for fermions each $q(j)$ and $p(j)$ can take only two different values ($\pm 1/2$), we see that the numerical computation (with a large but finite N) for fermions is much faster than that for bosons. Note

also that one can formally write (14) with the continuum notation as $Z = \int \mathcal{D}q(\tau) \mathcal{D}p(\tau) e^{-S}$, where $\tau = \beta j/N$,

$$\int \mathcal{D}q(\tau) \mathcal{D}p(\tau) \equiv \prod_{\tau=0}^{\beta} \frac{1}{2} \sum_{q(\tau)} \sum_{p(\tau)}, \quad (15)$$

$$S = \int_0^{\beta} d\tau \left[-i\pi \frac{dq(\tau)}{d\tau} p(\tau) + H(p(\tau), q(\tau)) \right], \quad (16)$$

but here the derivative $dq(\tau)/d\tau \equiv (q(j+1) - q(j))/\Delta\tau$ is not well defined because it is always equal to 0 or $\pm\infty$ as $\Delta\tau \rightarrow 0$. However, this is not a problem when S is calculated as in (13).

Let us now generalize the analysis to an arbitrary number of fermionic degrees of freedom. We label all degrees of freedom by $a = 1, \dots, L$, where a comprises various types of degrees of freedom, such as different space points, spinor indices, and color indices. The anticommutation relations are

$$\{\hat{\psi}_a, \hat{\psi}_b\} = \{\hat{\psi}_a^\dagger, \hat{\psi}_b^\dagger\} = 0, \quad \{\hat{\psi}_a, \hat{\psi}_b^\dagger\} = \delta_{ab}. \quad (17)$$

We can represent this algebra in terms of direct products of representations for each pair $\hat{\psi}_a, \hat{\psi}_a^\dagger$. The orthonormal basis for the 2^L -dimensional representation space can be chosen to be the set of 2^L states of the form

$$|n\rangle \equiv \bigotimes_a |n_a\rangle = |n_1\rangle \otimes \cdots \otimes |n_L\rangle, \quad (18)$$

where each n_a is equal to 0 or 1. We take the action of the operators on these states to be defined by

$$\hat{A}_1 \cdots \hat{A}_L |n_1\rangle \otimes \cdots \otimes |n_L\rangle = \hat{A}_1 |n_1\rangle \otimes \cdots \otimes \hat{A}_L |n_L\rangle, \quad (19)$$

where $\hat{A}_a = A_a(\hat{\psi}_a, \hat{\psi}_a^\dagger)$ with A_a being an arbitrary function. To determine the action of a differently ordered product of operators, one has to use the anticommutation relations (17) to write the action in terms of operators ordered as in (19). For example, $\hat{\psi}_2 \hat{\psi}_1 |n_1\rangle \otimes |n_2\rangle = -\hat{\psi}_1 \hat{\psi}_2 |n_1\rangle \otimes |n_2\rangle = -\hat{\psi}_1 |n_1\rangle \otimes \hat{\psi}_2 |n_2\rangle$. The hermitian conjugation of (19) determines the action of the operators on the left:

$$\langle n_L | \otimes \cdots \otimes \langle n_1 | \hat{A}_L^\dagger \cdots \hat{A}_1^\dagger = \langle n_L | \hat{A}_L^\dagger \otimes \cdots \otimes \langle n_1 | \hat{A}_1^\dagger. \quad (20)$$

To construct the functional formalism, we write $\hat{\psi}_a = \hat{q}_a + i\hat{p}_a$, so that $\hat{q}_a$ and $\hat{p}_a$ are hermitian operators that satisfy

$$\{\hat{q}_a, \hat{q}_b\} = \{\hat{p}_a, \hat{p}_b\} = (1/2)\delta_{ab}, \quad \{\hat{q}_a, \hat{p}_b\} = 0. \quad (21)$$

We introduce the states $|q\rangle = \bigotimes_a |q_a\rangle$ and $|p\rangle = \bigotimes_a |p_a\rangle$, which have the properties $\hat{q}_a |q\rangle = q_a |q\rangle$ and $\hat{p}_a |p\rangle = p_a |p\rangle$. These states satisfy the orthonormality and completeness relations analogous to (6) and (7), respectively, while the generalization of (8) is

$$\langle p | q \rangle = \frac{1}{\sqrt{2^L}} e^{-i\pi pq}, \quad (22)$$

where $pq \equiv \sum_a p_a q_a$.

We want to calculate the partition function for a Hamiltonian of the form

$$\hat{H} = \hat{\psi}^\dagger H^{(1)} \hat{\psi} + (\hat{\psi} H^{(2)} \hat{\psi} + \text{h.c.}), \quad (23)$$

where $\hat{\psi}^\dagger H^{(1)} \hat{\psi} \equiv \sum_{a,b} \hat{\psi}_a^\dagger H_{ab}^{(1)} \hat{\psi}_b$, and similarly for the second term. We can write

$$H_{ab}^{(1)} = A_{ab} + S_{ab}, \quad S_{ab} = S'_{ab} + \tilde{S}_{ab}, \quad (24)$$

where $A_{ab} = (H_{ab}^{(1)} - H_{ba}^{(1)})/2 = -A_{ba}$ is the antisymmetric part, $S_{ab} = (H_{ab}^{(1)} + H_{ba}^{(1)})/2 = S_{ba}$ is the symmetric part, $\tilde{S}_{ab} = \tilde{S}_a \delta_{ab}$, and $S'_{aa} = 0$. Using the anticommutation relations (21), we can write

$$\begin{aligned} \sum_{a,b} \hat{\psi}_a^\dagger H_{ab}^{(1)} \hat{\psi}_b &= (1/2) \sum_a \tilde{S}_a - 2i \sum_a \hat{p}_a \tilde{S}_a \hat{q}_a \\ &+ \sum_{a,b} \hat{q}_a A_{ab} \hat{q}_b + \sum_{a,b} \hat{p}_a A_{ab} \hat{p}_b. \end{aligned} \quad (25)$$

We want the $\hat{q}_a$ operators to be ordered as in (19), so we write

$$\sum_{a,b} \hat{q}_a A_{ab} \hat{q}_b = \sum_{a < b} \hat{q}_a A_{ab} \hat{q}_b + \sum_{a > b} \hat{q}_a A_{ab} \hat{q}_b$$

$$\begin{aligned} &= \sum_{a < b} \hat{q}_a A_{ab} \hat{q}_b + \sum_{a < b} \hat{q}_b A_{ba} \hat{q}_a \\ &= 2 \sum_{a < b} \hat{q}_a A_{ab} \hat{q}_b \\ &= 2 \sum_{a,b} \hat{q}_a A_{ab}^< \hat{q}_b, \end{aligned} \quad (26)$$

where

$$A_{ab}^< = \begin{cases} A_{ab}, & a < b, \\ 0, & a > b, \end{cases} \quad A_{ab}^> = \begin{cases} 0, & a < b, \\ A_{ab}, & a > b. \end{cases} \quad (27)$$

Similarly, we want the $\hat{p}_a$ operators to be ordered as in (20), so we write

$$\sum_{a,b} \hat{p}_a A_{ab} \hat{p}_b = 2 \sum_{a,b} \hat{p}_a A_{ab}^> \hat{p}_b. \quad (28)$$

We also apply a similar procedure for the second term in (23). Without losing on generality, the matrix $H^{(2)}$ is taken to be antisymmetric. Defining the matrices $A^{(+)}$ and $A^{(-)}$ as $A_{ab}^{(\pm)} = H_{ab}^{(2)} \pm H_{ab}^{(2)*}$, we can write the Hamiltonian (23) in the final form as

$$\begin{aligned} H(\hat{p}, \hat{q}) &= \frac{1}{2} \text{Tr} \tilde{S} - 2i \hat{p} \tilde{S} \hat{q} + 2 \hat{q} A^< \hat{q} + 2 \hat{p} A^> \hat{p} \\ &+ 2 \hat{q} A^{(-)<} \hat{q} - 2 \hat{p} A^{(-)>} \hat{p} + 2i \hat{p} A^{(+)} \hat{q}. \end{aligned} \quad (29)$$

This form of the Hamiltonian has the property

$$\langle p | H(\hat{p}, \hat{q}) | q \rangle = \langle p | q \rangle H(p, q). \quad (30)$$

Therefore, we can repeat the steps that led to (14) to obtain

$$Z = \lim_{N \rightarrow \infty} \left(\prod_{j=1}^N \frac{1}{2^L} \sum_{q(j)} \sum_{p(j)} \right) e^{-S}, \quad (31)$$

where S has the form (13) with the Hamiltonian $H(p, q)$ given by (29). In (31), $\sum_{q(j)} = \prod_a \sum_{q_a(j)}$, $\sum_{p(j)} = \prod_a \sum_{p_a(j)}$.

Needless to say, the formalism developed above can also be applied to the calculation of the partition function at finite chemical potential. Moreover, essentially the same formalism can be applied to the calculation of correlation functions among fields at different times, in a way similar to that for bosons. Using the fact that $\hat{H}$ is the generator of infinitesimal time translations, one finds similar expressions with $\Delta\tau \rightarrow i\Delta t$.

Concerning the application of our formalism to concrete physical models, two additional remarks may be in order. First, one should be careful when the Hamiltonian is written in terms of Dirac fields $\hat{\psi}^D, \hat{\psi}^{D\dagger}$. For example, if there are two degrees of freedom, one corresponding to a particle and the other to an antiparticle, then the ‘‘Dirac field’’ is $\hat{\psi}^D = (\hat{\psi}_1 + \hat{\psi}_2^\dagger)/\sqrt{2}$. Second, although the operator $\hat{N} = \sum_a \hat{\psi}_a^\dagger \hat{\psi}_a$ has the property

$\hat{N}|n\rangle = (\sum_a n_a)|n\rangle$, the operator of the physical number of particles, in general, has a more complicated form $\hat{N}_{\text{phys}} = \sum_{a,b} \hat{\psi}_a^\dagger N_{ab} \hat{\psi}_b$. For the explicit form of N_{ab} in the continuum case, see [10].

In this Letter, we have presented a general formalism rather than apply it to concrete physical problems. However, it is straightforward to apply the developed formalism to any physical model, such as lattice QCD. We strongly believe that this formalism will be widely applied to lattice QCD and lead to a number of important phenomenological results that previously could not be obtained because they required too much computer time with previous algorithms for fermions.

Finally, let us shortly discuss the representation of quantum fermionic states in terms of wave functions $\Psi(q)$ written in the q -representation. Note that it is already known how to represent fermionic quantum states in terms of wave functions $\Psi(\psi)$, where ψ is a Grassmann variable [11]. However, $\Psi(\psi)$ is a Grassmann valued quantity, so $\Psi^*(\psi)\Psi(\psi)$ cannot be interpreted as a probability density. This problem does not exist for $\Psi(q)$. We first introduce the base functions

$$\Psi_+(q) = \langle q|q_+\rangle = \Theta(q), \quad \Psi_-(q) = \langle q|q_-\rangle = \Theta(-q), \quad (32)$$

where $\Theta(q)$ is the step function equal to 1 (0) for $q = 1/2$ ($q = -1/2$). From (3) and (5) we see that $\hat{p}|q_\pm\rangle = \mp(i/2)|q_\mp\rangle$. Therefore, we represent the operators $\hat{q}$ and $\hat{p}$ as

$$\hat{q} = q, \quad \hat{p} = -iq\hat{e}_q, \quad (33)$$

where $\hat{e}_q$ is the operator that acts on the space of functions as $\hat{e}_q f(q) = f(-q)$. We see that the operator $q\hat{e}_q$ plays the same role for fermions as the derivative operator ∂_q does for bosons. The operators (2) are now represented as

$$\hat{\psi} = q(1 + \hat{e}_q), \quad \hat{\psi}^\dagger = q(1 - \hat{e}_q). \quad (34)$$

It is easy to show that, for example, $\{\hat{p}, \hat{p}\}f(q) = 2q^2 f(q)$, $\{\hat{\psi}, \hat{\psi}^\dagger\}f(q) = 4q^2 f(q)$, which is in agreement with the

anticommutation relations (4) and (1), having in mind that the operators act on functions $f(q)$ defined for $q \in \{-1/2, 1/2\}$. For time dependent wave functions, i.e., in the Schrödinger picture, the Schrödinger equation is

$$H(-iq\hat{e}_q, q)\Psi(q, t) = i\partial_t\Psi(q, t). \quad (35)$$

The expected value of an arbitrary operator $\hat{O} = O(\hat{q}, \hat{p})$ is given by

$$\langle\hat{O}(t)\rangle = \sum_q \Psi^*(q, t)O(q, -iq\hat{e}_q)\Psi(q, t). \quad (36)$$

The generalization of Eqs. (32)-(36) to the case with many fermionic degrees of freedom is obvious.

The author is grateful to N. Bilić and H. Štefančić for critical reading of the manuscript. This work was supported by the Ministry of Science and Technology of the Republic of Croatia under Contract No. 0098002.

* Electronic address: hrvoje@thphys.irb.hr

- [1] F. A. Berezin, *The method of second quantization* (Academic Press, NY, 1966).
- [2] C. Davies, Lectures given at 55th Scottish Universities Summer School, St. Andrews, August 2001; hep-ph/0205181.
- [3] S. Duane, A. D. Kennedy, B. J. Pendleton, and D. Roweth, Phys. Lett. B **195**, 216 (1987).
- [4] M. Lüscher, Nucl. Phys. B **418**, 637 (1994).
- [5] A. A. Slavnov, Phys. Lett. B **388**, 147 (1996).
- [6] M. Creutz, Phys. Rev. Lett. **81**, 3555 (1998).
- [7] T. Bakeyev, Phys. Lett. B **519**, 277 (2001).
- [8] M. Creutz, Nucl. Phys. Proc. Suppl. **94**, 219 (2001).
- [9] J. I. Kapusta, *Finite-temperature field theory* (Cambridge University Press, Cambridge, 1989).
- [10] H. Nikolić, Phys. Lett. B **527**, 119 (2002); Erratum-ibid. B **529**, 265 (2002); hep-th/0202204, to appear in Int. J. Mod. Phys. D; hep-th/0205022.
- [11] R. Floreanini and R. Jackiw, Phys. Rev. D **37**, 2206 (1988); J. Hallin and P. Liljenberg, Phys. Rev. D **52**, 1150 (1995).